AUTHOR INDEX

Ahmed, I. A., *see* Alim, K. A.
Akehurst, B. C., Turkish tobacco in Southern Rhodesia, 267–80.
Alim, K. A., month, age, and intervals of calving in Egyptian dairy buffaloes, 37–41.
Ashfaq, M., environment and heredity in Pakistani buffalo milk yield, 161–75.
Ashton, E. D., milk recording and use of milk records for stock improvement, 1–9.
Asker, A. A., effect of crossing on growth of Egyptian lambs, 256–60.
Asker, A. A., *see* Ragab, M. T.

Barrowman, J. R., the rubber-ring ligature for castrating and docking lambs, 189–202.
Bastawisy, A. E., *see* Asker, A. A.
Boaz, T. G., *see* Barrowman, J. R.
Bogdan, A., control of Aristida in Kenya Rift Valley pastures, 211–23.
Brown, N. J., two soil tests for cultivation investigations, 138–40.

Chenery, E. M., *see* Pereira, H. C.
Coles, R., hatching rates in hens' eggs, 281–92.
Crowther, E. M., *see* Haines, W. B.

Durham, V. M., *see* Selman, I. W.

El-Issawi, H. F., *see* Khishin, S. S.

Fountaine, E. R., *see* Brown, N. J.

Gordon, J., uniformity trial on Gold Coast cocoa, 332.

Haines, W. B., manuring Hevea. V. Long-term effects in Malaya, 203–10.
Hammond, G. H., fluctuations in white-grub populations in Ontario, 59–64.
Hattingh, E. R., control of *Stoebe vulgaris* by selective weedkillers, 10–18.
Hayward, L. A. W., infestation of cocoa by tobacco beetle in Nigeria, 133–7.

Jones, P. A., *see* Pereira, H. C.

Khishin, S. S., the Jersey in Egypt, 121–7.
Knight, R. L., cotton breeding in the Sudan. I. Egyptian cotton, 68–80. II. Egyptian cotton (*continued*), 81–92. III. American cotton, 176–84.

Large, R. V., the growth of Clun lambs, 141–7.

Mahadevan, P., heritability of milk yield in Indian-European dairy crosses, 93–96.
Mason, I. L., *see* Ashfaq, M.
Mills, W. R., *see* Pereira, H. C.
Moore, R. M., control of hoary cress (*Cardaria draba*) by phenoxyacetic derivatives, 112–20.

Nye, P. H., value of fertilizers in the Gold Coast. III. Voltaian Sandstone and southern maize areas, 42–54.
Nye, P. H., Gold Coast fertilizer responses: effects of time and method of application, 101–11.

Payne, P. C. J., *see* Brown, N. J.
Pereira, H. C., coffee responses to manuring and mulching, 23–36.
Pereira, H. C., tillage of Kenya coffee. I. Yields, 231–40. II. Soil structure, 323–31.
Pereira, H. C., transient effects of grasses on tropical soil structure, 148–60.

Ragab, M. T., twinning in Ossimi sheep, 224–30.
Ragab, M. T., *see* Asker, A. A.
Reith, J. W. S., effects of calcic and magnesian liming materials on crop composition, 305–13.
Ripley, P. O., weed control in western Canada, 261–6.
Russell, T. A., potato blight in West Africa, 19–22.

Sabet, K. A., new bacterial disease of maize in Egypt, 65–67.
Selman, I. W., organic and inorganic fertilizers on potatoes, 314–22.
Sheffield, F. M. L., erinose of sweet potato, 97–100.
Shepherd, R. W., use of nitrogen for sugar-beet on light land, 128–32.
Spedding, C. R. W., persistence of worm-infestation effects in sheep, 55–58.
Storrar, A., *see* Bogdan, A.

Tayler, J. C., *see* Large, R. V.
Thomson, J. R., effect of seed dressings on oat germination tests, 185–8.
Thornton, G. J., *see* Haines, W. B.
Towers, K. G., *see* Barrowman, J. R.

Underwood, J. R., *see* Coles, R.

Walker, P. T., influence of climate on wheat aphid outbreak in Kenya, 293–304.
Wyllie, J., wartime food production in the U.K., 241–55.

SUBJECT INDEX

Aphid, wheat, influence of climate on, in Kenya, 293–304; biology, aphid, in relation to climate, 300–2; control, natural and by spraying, 303; meteorological conditions, 294–300; outbreaks, aphid, severity and area, 293–4, 296; population dynamics, 302–3; rainfall distribution, monthly, 296–8; temperatures, related to outbreaks, 298–300.
Aristida, control of, in Kenya pastures, 211–

23; annual grasses, occurrence of, 211–12; burning, effects of, 219–20; control secured by burning or heavy seasonal grazing, 221–2; grazing by cattle or sheep, and effects on herbage, 213–22; grazing preferences, 215; mowing, effects of, 220; rainfall, monthly, 213; seasonal grazing and effects, 213–22.

Beet, sugar, *see* Sugar-beet.
Buffalo, Egyptian dairy, calving in, 37–41; age at first calving, 39; calving intervals and lactation periods, 40–41; month of calving, 37–38.
Buffalo, Pakistani, milk yield of, 161–75; age at calving, effect on milk yield, 169–71; age at first calving, 165; breeding policy recommended, 174; breeds of buffalo in west Pakistan, 161; calving interval, 166–7; environmental and genetical effects on milk yield, 161–74; Government Buffalo Breeding Farm, 162–3; heritability of milk yield, 172–4; lactation records studied, 163–4; length of lactation, 167–8; month of calving, 165–6; repeatability of milk yield, 171–2.

Cacao, Gold Coast, uniformity trial on, 332; optimum tree density, 332.
Calcium and magnesium contents of crops, *see* Liming materials, calcic and magnesian.
Cardaria draba, *see* Hoary cress, control of.
Cattle, dairy, milk yield in crosses between Indian and European breeds, 93–96; cross-bred cattle, milk production of, 94; heritability of milk yield, 95–96; milk records studied, 93–94; repeatability of milk yield, 94–96.
Cattle, dairy, *see* Jersey, the, in Egypt.
Cocoa, infestations of, by tobacco beetle in Nigeria, 133–7; climatic preferences of tobacco beetle, 134–6; cowpea trade, relation to cocoa infestation, 134–6; *Lasioderma serricorne*, damage by, 133–4; origins of infestation of cocoa, 134–6.
Coffee, Kenya, field responses to fertilizers, manures, and mulches, 23–36; cattle manure, effect of, 27, 31; field experiments, modern, 24–35; field trials, early, 24–26; grass mulching and its problems, 31–35; mulches, grass, effect on soil structure, 33–34; mulches, grass, effect on yields, 32–33; nitrogen, phosphate, and potash responses, 27–31; nutrient requirements of coffee, 23–24; phosphate status of soils, 28–29; phosphate uptake of coffee, 28.
Coffee, Kenya, tillage of. I. Effects on coffee yields, 231–40; cultivation, avoidance of, 239; current practice, improvements in, 238–9; depth of tillage, effect on yield, 235–7; die-back, effect of tillage on, 235; erosion, subsoiling, and weed-growth problems, 231; implement type, effect on yield, 235; manure, cattle, response to, 237; mulch, grass, use of, 239; quality of coffee, effect of tillage on, 238; weeds, serious effect on yields, 233–5. II. Effects on soil structure, 323–31; aggregation, soil, effects of implements on, 327–8; clean-weeding, effects on structure, 324–6; field observations of tilth, 323; pore space, soil, effects of implements on, 327; rotary cultivator, improvement of, 329–30; volumetric structure measurements contrasted with wet-sieving, 328–9.
Compost, effects of, *see* Potatoes, growth and yield of.
Cotton breeding in the Sudan. I. Egyptian cotton, 68–80; blackarm disease and resistant varieties, 71–75; control of blackarm by seed disinfection and agricultural measures, 76–77; damage caused by diseases and pests, 70–71; evolution of blackarm resistance, 75–76; leafcurl resistance, 77–79; types of cotton grown, 69–70. II. Egyptian cotton (*continued*), 81–92; bollworm resistance, 90–91; cluster and short-branch characters, 88–89; deciduous bracts, 85–86; earliness, 89–90; grade, association with blackarm resistance, 83–84; jassid resistance, 81–83; locules per boll, ovules per locule, 88; salt tolerance, 84–85; storm resistance, 86; strength of lint, 86–87; yield potential, 87. III. American upland cotton, 176–84; blackarm resistance, 176–9; breeding methods used in the Sudan, 181–2; cluster types, 180; deciduous bracts, 179–80; earliness and pest avoidance, 181; ginning out-turn, 180; varieties, Sudan, key to, 183–4.
Crosses between Indian and European breeds, *see* Cattle, dairy, milk yield in.
Crosses between Jersey and Egyptian cattle, *see* Jersey, the, in Egypt.
Crossing of lambs, effect on growth, *see* Lambs, Egyptian.
Cultivation investigations, two soil tests for, 138–40; aggregate analyser, mechanical, 138–9; volume weight, determination by excavation, 139–40.

Dairy cattle, *see* Cattle, dairy, milk yield in; Jersey, the, in Egypt. *See also* Buffalo, Egyptian dairy; Buffalo, Pakistani, milk yield of.
Dairy stock improvement, use of milk records in, 1–9; breed structure of the national herd, 4–5; bull selection and use of progeny tests, 5–7; milk recording in England and Wales, 1–4; other uses of milk records, 7–8.

Eggs, hens', hatching in, *see* Hatching rates.
Erinose of sweet potato, *see* Potato, sweet.

Fertilizer responses in the Gold Coast, relation to time and method of application, 101–11; ammonium sulphate, residual effects of, 108–9; fertilizers, qualities of, 105; mixed crops, experiments with, 106–7; nitrogen and phosphate, effects on mixed crops, 106–7; phosphate, rates, times, and methods of application of, 101–4, 106–7; placement of phosphate, effects

of, 103–4; superphosphate, residual effects of, 108–10.
Fertilizers, survey of the value of, in Gold Coast food-farming areas. III. Voltaian Sandstone and southern maize areas, 42–54; agriculture, local, 44, 45, 51; climate, 43, 50; cropping history and fertilizer responses, 46–47; nitrogen and phosphate, responses to, 45–49, 52–53; southern maize area, value of fertilizers to, 52–53; Voltaian Sandstone area, best use of fertilizers in, 48–49.
Food, human, wartime production of, in the U.K., 241–55; acreage of tillage land, importance of, 253; acres to feed one person, 248–9; annual index of calorie production, 250; calorie output, percentage composition of, 247–8; calorie production per acre, 243–4; per man engaged, 252–3; calorie requirements per head, 241–2; changes in sources of calories, 244–6; crops, areas and yields of, 246, 253; imports of feedingstuffs, 251–2, 254; increased food production, factors in, 253–5; livestock, 246–7; persons, numbers of, fed from home production, 249–50; price per million calories, 250–1; wartime compared with pre- and post-war food production, 244–51, 253–5; yields per acre, 246, 253.

Germination of oats, effects of seed dressings on, *see* Seed dressings.
Grasses, transient effects of, on soil structure, *see* Structure of tropical soils.
Grubs, white, in sod, *see* White grubs (*Phyllophaga* spp.).

Hatching rates in hens' eggs, observations on, 281–92; breeds, differing performance of, 288–9; disease, influence on hatching rates, 285–6; egg weight, selection based on, 290; factors lowering hatchability, 282–6; losses, serious, from low hatching rates, 281–2; management, faulty, 283; nutrition, importance of, 284; results, limited, from experiments to improve hatching, 287–91.
Herbicides, selective, *see* Hoary cress (*Cardaria draba*); *Stoebe vulgaris*; Weed control in western Canada.
Hevea, manuring. V. Long-term effects in Malaya, 203–10; edge effects in plots, 203–4; growth increases after manuring ceased, 204–8, 209–10; nitrogen and phosphates, residual effects of, 205–10; phosphate, high economic return from, 208; 'poaching' of fertilizer treatments, 203–4; yield results, 207–9.
Hoary cress (*Cardaria draba*), control of, by chlorinated phenoxyacetic compounds, 112–20; concentration, effect of, 113–15, 117; dosage response curve, 114; double and single applications, comparison of, 115–19; MCPA, sodium salt, control by, 113–19; seasonal fluctuations in populations of Cardaria, 114–16; 2,4-D, salts and esters of, compared, 117–19; 2,4,5-T amine, 117, 119.

Jersey, the, in Egypt, 121–7; age at first calving, 125; calving intervals, 124; cross- and pure-bred Jersey cows compared, 122–6; heat tolerance of Jerseys, 123; introduction of Jersey cattle, 121; length of lactation, 123–4; milk yields from Jersey cows and crosses with Egyptian cattle, 122–3; other breeds compared with Jerseys in Egypt, 122–3.

Lambs, castration and docking of, rubber-ring technique for, 189–202; advantages and disadvantages of rubber-ring method, 189, 201; age, best, for castration and docking, 200–1; castrating, physical effects of, 189, 192–3, 195–7, 199–201; docking, physical effects of, 189–90, 193–5, 197–201; early treatment preferable, 200–1; Elastrator (rubber-ring ligature), use of, 189–201; weight performance of treated animals, 195–9.
Lambs, Clun, studies in the growth of, 141–7; birth weights and measurements, 142–4; ewes and rams compared, 143–8; management and nutrition, 141–2; measurements, useful, to indicate plane of nutrition, 146–7; nutrition, high and low, effects on weight and size, 144–7; singles and twins compared, 143–4, 146; skeletal growth, 145–7; weight increases, 144–5.
Lambs, Egyptian, effects of crossing on growth of, 256–60; cross-bred lambs, superiority of, 258–9; daily gains in weights and measurements, 258–9; measurements, average, 257–9; mutton production, value of cross-breeding for, 259; Ossimi and Rahmani lambs and crosses, 256–9; weights, average, 257–9.
Liming materials, calcic and magnesian, effects on crop composition, 305–13; arable crops, calcium and magnesium contents of, 307–9, 312; grassland herbage, magnesium content of, increased by magnesian limes, 308–12; seasonal variations in composition of Scottish grassland, 309–12; soils, composition of, after liming treatments, 311–12; yields of crops and herbage, effects of liming materials on, 307–10, 312.

Maize, new bacterial disease of, in Egypt, 65–67; *Erwinia carotovora* f.sp. *zeae*, proposed as name, 67; pathogen, description of, 66; soft-rot symptoms observed, 65.
Manures and fertilizers, effects of: *see* Coffee, Kenya, field responses; Fertilizer responses, and fertilizer survey, in the Gold Coast; Hevea, manuring; Potatoes, growth and yield of; Sugar-beet, use of nitrogen; Tobacco, Turkish or oriental.
Milk recording in England and Wales, *see* Dairy stock improvement, use of milk records in.
Milk yield, environmental and genetical effects on, *see* Buffalo, Pakistani.
Milk yield in Indian-European crosses, *see* Cattle, dairy.
Milk yield in Jersey-Egyptian crosses, *see* Jersey, the, in Egypt.

Mulches, grass, effects of, *see* Coffee, Kenya, field responses.

Nitrogen, use of, *see* Sugar-beet.

Oats, germination tests, effect of seed dressings on, *see* Seed dressings.
Organic and inorganic fertilizers, comparison of, *see* Potatoes, growth and yield of.
Organo-mercurial and gamma BHC seed dressings, effect on germination, *see* Seed dressings.

Pastures, Kenya, control of annuals in, *see* Aristida.
Phenoxyacetic derivatives in weed control, *see* Herbicides, selective.
Phyllophaga spp., *see* White grubs.
Population fluctuations, long-term, *see* White grubs.
Potato blight in W. Africa, 19–22; occurrence in British Cameroons, 19–20; resistance of imported potato varieties, 21.
Potato, sweet, erinose of, 97–100; control of disease, 98–100; distribution in Africa, 97, 100; downy or hairy appearance of plants, 97–99; erinose or hairiness, due to mites, 97–98; yield, effect of disease on, 99–100.
Potatoes, growth and yield of, effects of manures, 314–22; compost, comparative effects of, on plant growth, 317–19, 321, on tuber yield and composition, 320–2; fertilizers, chemical, compared with organics, 317–22; height, as affected by manuring, 317, 321; nitrogen content of plants and tubers, 318–21.

Rubber, manuring, *see* Hevea.
Rubber-ring castration and docking of lambs, *see* Lambs, castration.

Seed dressing, effect on germination tests of oats, 185–8; abnormal seedlings produced, 185–7; gamma BHC and organo-mercurial dressing, slight effects of, 185–7; tests, germination, on sand or soil, 186–7.
Sheep, Ossimi, twinning in, 224–30; age, effect of, on lambing percentage, 225–6, 228; birth, type, influence of, 226; heritability of twinning, 227, 229; inbreeding, effect of, 226, 228; mating time, effect of, 225, 228; ram or sire, influence of, 226–9; year, influence of, 224–5, 228.
Sheep, persistence of effects of worm infestation in, 55–58; carcass quality, effect of infestation on, 57; liveweight, effect on, 55–57; resistance, age, development of, 57–58; wool production, effect on, 57.
Soil structure, *see* Coffee, Kenya, tillage of; Structure of tropical soils.
Soil tests for cultivation investigations, *see* Cultivation investigations.
Stoebe vulgaris, control of, by selective weedkillers, 10–18; aerial spraying, success of, 16; burning, influence on herbicide action, 12–14, 16–17; management and mowing, influence of, 11–17; phenoxyacetic derivatives, comparison of salts and esters of, 10–17; root reserves, effect of herbicides on, 14.
Structure of tropical soils, transient effects of grasses on, 148–60; aeration and pore-space, influence of cropping on, 154–6; compactness, soil, 153; crops compared, cover, cultivated and grass, 149; deterioration of structure, rate of, 156–9; grasses, physical effects of, better than cover crops, 159; infiltration and percolation, 155–6; porosity, soil, 153–5; rainfall, artificial, value of technique, 156, 159; rotations, crop, comparison of effects on structure, 149–60; sieving, wet, techniques of, 149–53.
Structure, soil, *see* Coffee, Kenya, tillage of. II.
Sugar-beet, use of nitrogen for, on light land, 128–32; beet, clean, yield of, 128–30; optimum nitrogen dressing, 132; purity of juice, 130–1; rainfall, relation of nitrogen response to, 131–2; sugar percentage, 130; sugar, yield of, 130; tops, yield of, 131.

Tillage, effects on coffee yields, *see* Coffee, Kenya, tillage of. I.
Tillage, effects on soil structure, *see* Coffee, Kenya, tillage of. II.
Tobacco beetle, infestation of cocoa by, *see* Cocoa, infestations.
Tobacco, Turkish or oriental, in Southern Rhodesia, 267–80; fertilizers, effects on seedling numbers, 274, on the crop, 275–80; grade, fertilizer effects on, 275–9; nitrogen, phosphate, and potash, effects on yield and quality, 276–80; planting, best time of, 274; rates of fertilizers, optimum, 279–80; seeding rates, 273; seedlings, production of, 272–4; spacing, 275; varieties, characteristics and comparison of, 268–72; weed control in seed-beds, 273.
Twinning, in sheep, *see* Sheep, Ossimi.

Weed control in western Canada, 261–6; areas sprayed with 2,4-D, 264; chemicals, early use of, 262–3; cropping and tillage methods, 261–2; growth regulators, effectiveness and rapid development of, 263–5; herbicides, selective, 262–5; problem, weed, the, 261; 2,4-D, development and use of, 264–5.
Weed control, *see* Aristida; Herbicides, selective.
Wheat aphid in Kenya, influence of climate on, *see* Aphid, wheat.
White grubs, population fluctuations of, in sod, 59–64; brood areas, in Ontario, 59; cycle, three-year, in *Phyllophaga* spp., 60–63; populations, annual, and percentage changes, 62–63.
Worm infestation, persistence, *see* Sheep, persistence of effects.

PRINTED IN GREAT BRITAIN